Navigating Relational Transitions and Antidepressant-Induced Emotional Blunting

Navigating Relational Transitions and Antidepressant-Induced Emotional Blunting

Sara Larson

Copyright © 2026 by Sara Larson
All rights reserved. No part of this book may be reproduced in any manner whatsoever without written permission except in the case of brief quotations embodied in critical articles and reviews.
First Printing, 2026

Introduction

the Intersection of Psychopharmacology and Relationship Dynamics

The pharmacological management of major depressive disorder (MDD) and severe anxiety disorders relies heavily on the administration of Selective Serotonin Reuptake Inhibitors (SSRIs) and Serotonin-Norepinephrine Reuptake Inhibitors (SNRIs). These medications are foundational in modern psychiatry, frequently providing necessary mood stabilization and crisis prevention for millions of patients globally. However, the introduction of these powerful neuromodulators carries complex, systemic side effects that extend far beyond the individual patient, often inducing profound disruptions within the patient's interpersonal ecosystem. Among the most pervasive and relationally destructive of these secondary effects is Antidepressant-Induced Emotional Blunting (AIEB), a clinical phenomenon characterized by the pervasive dampening of the patient's entire emotional spectrum.[1]

When a patient experiencing AIEB initiates the de-escalation of a romantic relationship—frequently manifesting as a request to "just be friends"—the non-depressed partner is thrust into a highly destabilizing state of emotional and psychological limbo. Clinical observations suggest that this request is rarely rooted in malice or organic incompatibility. Rather, it is a direct byproduct of the psychological flattening, apathy, and loss of reinforcement learning fundamentally in-

duced by the serotonergic medication.[3] The depressed patient, increasingly unable to feel or process the passionate, high-valence emotions required to sustain romantic and physical intimacy, naturally seeks to downgrade the relationship to a level that matches their artificially restricted emotional bandwidth.[4]

The transition from a romantic partnership to a platonic friendship under the overarching shadow of antidepressant-induced emotional blunting requires careful clinical observation, rigid boundary setting, and a deep understanding of psychological grief models such as ambiguous loss. If this transition is mismanaged, the healthy partner risks falling into a state of chronic caregiver burnout, while the depressed partner may inadvertently utilize the resulting "friendship" as a buffer to avoid addressing the adverse neurochemical effects of their psychiatric treatment.[5] The comprehensive analysis that follows explores the neurobiological origins of this pharmacological blunting, the severe psychological toll exacted on the healthy partner, the vital mechanisms of boundary realignment, and the specific clinical pathways available for couples navigating this profound relational crisis.

The Neurobiology and Clinical Reality of Emotional Blunting

To effectively navigate the relational fallout of an SSRI-induced breakup or transition, it is imperative to first understand the neurobiological mechanisms driving the depressed partner's altered behavior. Antidepressant-induced emotional blunting is not a rare or isolated occurrence; empirical data in-

dicates that between 40% and 60% of patients taking SSRIs or SNRIs experience some degree of emotional dampening, with some clinical surveys suggesting the prevalence may be even higher.[1]

The Mechanism of Serotonin and Dopamine Imbalance

Selective Serotonin Reuptake Inhibitors function by inhibiting the reuptake of serotonin in the synaptic cleft, thereby increasing the availability of this neurotransmitter to alleviate the crushing lows, rumination, and despair associated with clinical depression.[2] However, the human brain operates as a deeply interconnected and highly sensitive neurochemical ecosystem. The artificial influx and sustained pooling of serotonin can inadvertently suppress dopamine transmission, particularly within the brain's reward pathways.[1] Dopamine is the primary neurotransmitter responsible for motivation, pleasure, anticipation, and the high-intensity feelings typically associated with romantic love and sexual desire.

Recent neurobiological studies have successfully isolated the specific cognitive functions impaired by chronic SSRI administration. A landmark clinical study conducted by researchers at the University of Cambridge, in collaboration with the University of Copenhagen, recruited 66 healthy volunteers to investigate these mechanisms.[3] The cohort was divided, with half receiving the well-tolerated SSRI escitalopram and the other half receiving a placebo over a minimum of 21 days. The researchers discovered that while overall cognition remained intact, the SSRI significantly impaired "reinforcement learning"—the critical behavioral process by

which humans learn from and respond to positive environmental feedback.[3] In a relational context, this impairment means that the patient literally loses the neurological ability to experience joy, reward, or positive reinforcement from everyday social interactions, including the presence, affection, or support of a romantic partner.

Affective to Cognitive Processing Shifts

A hallmark of AIEB is a fundamental shift in how the patient interacts with the world, moving from a fully integrated emotional experience to a detached, observational state. Clinical observations note a distinct transition from affective (feeling-based) processing to cognitive (thought-based) processing.[8] The patient may cognitively recognize that they care deeply for their partner, but they find themselves entirely disconnected from the somatic and affective experience of that love.

This processing shift creates a chilling dynamic in romantic partnerships. The depressed partner may articulate that they "know" they love their significant other, but they "feel" absolutely nothing.[2] This state of emotional detachment leads to a pervasive feeling of being a passive observer in one's own life. The patient is stripped of enthusiasm, passion, and affection, while simultaneously losing access to appropriate negative emotions, such as the ability to process grief, worry, or anger when circumstances warrant it.[8] Some literature unifies this overarching state under the moniker "selective serotonin reuptake inhibitor-induced indifference," highlighting the insidious onset and complete resolution of symptoms only

upon the discontinuation or adjustment of the offending drug.[9]

Differentiating Antidepressant-Induced Blunting from Depressive Anhedonia

A critical clinical challenge in managing these relationship transitions is determining whether the partner's emotional withdrawal is a symptom of residual, untreated depression (anhedonia) or an iatrogenic side effect of the medication (AIEB). Recognizing this distinction is paramount for the non-depressed partner, as it dictates the reality of the relational collapse. If the apathy is drug-induced, the loss of romantic connection is a pharmacological artifact, not necessarily an organic evolution of the couple's compatibility.[4]

Diagnostic Domain	Depressive Anhedonia	En
Onset and Chronology	Precedes psychiatric treatment; constitutes a core feature of the initial major depressive episode.	*af* cr tic
Emotional Valence Profile	Primarily targets and eliminates positive emotions (inability to feel pleasure or joy); negative emotions (sadness, despair, psychic pain) remain intensely active and painful.	tiv tie or se sti

Cognitive and Psychological State	Often accompanied by intense, active rumination, overwhelming guilt, and profound feelings of worthlessness.[10]	in ca pu pr
Physiological Comorbidities	Frequently accompanied by severe lethargy, significant sleep disturbances, and active hopelessness.[10]	qu en ov su

When a partner on SSRIs requests to transition the relationship to a friendship, they are often attempting to reconcile their cognitive memory of the relationship's value with their current inability to generate the affective responses required to sustain it. The patient experiences a reduction in sympathy and empathy, reporting that feelings toward partners and children are disconnected or detached from their own instincts.[11]

The Psychological Trauma and Impact on the Non-Depressed Partner

When a romantic relationship is subjected to the prolonged effects of AIEB, the non-depressed partner endures a highly specific and deeply disorienting form of psychological trauma. The request to remain friends while simultaneously stripping away the romantic, passionate, and intimate components of the relationship creates profound cognitive dis-

sonance. The healthy partner is forced to interact with the physical embodiment of their significant other while being starved of the emotional reciprocity that defines the bond.

The Dynamics of Ambiguous Loss

The theoretical framework of "ambiguous loss" is highly applicable to relationships affected by SSRI blunting. Ambiguous loss occurs when a person is physically present but psychologically or emotionally absent.[5] The non-depressed partner is forced to mourn the loss of the vibrant, emotionally available person they once knew, even while that person continues to sleep in the same bed or text them daily.

Transitioning immediately into a platonic friendship exacerbates this ambiguous loss. The continued physical proximity and ongoing communication serve to keep the healthy partner's attachment system highly activated, while the depressed partner's emotional flatness constantly starves that system of validation and reciprocity.[5] The emotional numbness associated with SSRIs is rarely dramatic; it operates as a quiet killer, eroding the foundation of the relationship slowly as conversations shift from deep vulnerability to shallow, logistical check-ins.[5] The healthy partner inevitably interprets this mechanical intimacy and lack of spark as a personal failure or definitive evidence that they are unlovable, rather than accurately recognizing it as a neurochemical deficit within their partner.[5]

The Illusion of Friendship and the Disruption of Self-Expansion

Close relationships rely heavily on the psychological concept of "self-expansion," wherein partners incorporate aspects, experiences, and perspectives of one another into their own identities. This cognitive reorganization of the self is highly correlated with improved mental health, and research utilizing cross-sectional, daily diary, and longitudinal methodologies demonstrates that greater self-expansion is robustly associated with fewer depressive symptoms.[16]

When one partner becomes emotionally numb due to pharmacological interventions, the avenue for mutual self-expansion abruptly closes. The healthy partner's identity may begin to erode, systematically replaced by the singular, demanding role of the "supportive friend" or "crisis manager." The preservation of the healthy partner's distinct psychological identity is severely compromised when they agree to a nebulous, undefined friendship with a blunted ex-partner.[16] The relationship enters a state of persistent emotional numbness, leading to acute isolation, loneliness, and resentment for the non-depressed partner.[14]

Caregiver Burnout in Romantic Transitions

When a partner suffering from AIEB suggests friendship, the request is often a subconscious psychological strategy to maintain the stabilizing presence of the healthy partner without having to meet the emotional obligations of a romantic relationship.[5] The depressed partner heavily benefits from the predictability, routine, and practical support the relationship provides, but this arrangement is inherently asymmetrical.

By accepting the "friendship" without enforcing strict boundaries, the non-depressed partner frequently slips into the role of an unpaid clinical caregiver. The psychological literature on caregiving within romantic frameworks highlights that assuming a caretaking role strips away romantic intimacy and drastically increases the risk of depression, social isolation, and burnout in the healthy partner.[7] Caregivers already exhibit higher rates of stress and social isolation than non-caregivers, and the emotional exhaustion inherent in balancing caregiving with personal relationship expectations fundamentally alters the dynamic.[19]

The healthy partner may rationalize the situation, believing that by remaining close friends, they can continuously monitor the depressed partner's recovery, ensure medication compliance, and eventually "win them back" once the psychiatric medication is successfully adjusted. This hope is functionally maladaptive; it delays the healthy partner's own psychological healing, traps them in an indefinite holding pattern, and fosters deep-seated resentment.[19] The internal conflict arises when the healthy partner's logical voice—which demands self-preservation—is challenged by their internal "rescuer," which questions their compassion if they are not constantly sacrificing their own needs for the sick partner.[7]

Clinical Strategies for the Transition: The Imperative of Boundaries

If a couple decides to attempt a transition from romantic partners to friends while one partner navigates complex SSRI treatment, standard breakup advice is entirely insufficient.

The presence of a severe psychiatric variable requires structured, clinical-grade boundary setting to protect the mental health of the non-depressed individual, prevent caregiver burnout, and force the depressed individual to confront the reality of their pharmacological state.[22]

The Necessity of a No-Contact Period

An immediate pivot from lovers to friends is psychologically untenable, particularly when one partner is suffering from medication-induced apathy. A mandatory "no-contact" period is a critical first step in restructuring the relationship.[22] This period serves multiple vital functions:

1. **Detoxification of the Attachment System:** It allows the healthy partner's nervous system to deregulate from the constant, exhausting stress of seeking emotional validation from an unresponsive, chemically blunted source.[25]
2. **Assessment of Baseline Reality:** It forces the depressed partner to experience daily life without the buffering, regulating presence of their ex-partner. This deprivation may provide the necessary emotional friction for the depressed partner to recognize the severity of their emotional blunting and subsequently address it with their psychiatric provider.[2]
3. **Establishment of New Norms:** It clearly demarcates the definitive end of the romantic dynamic, ensuring that when contact eventually resumes, it is built on an entirely new foundation rather than serving as a diluted continuation of old, toxic habits.[22]

Boundary Matrix: Restructuring the Relationship Ecosystem

Once contact resumes after a cooling-off period, the parameters of the friendship must be explicitly defined and rigidly enforced. Boundaries must shift permanently from those characteristic of an intimate partnership to those of a standard, platonic friendship.[23] Boundary setting in recovery contexts involves physical, emotional, temporal, and digital limits.[23]

Boundary Domain	Romantic Partnership (Pre-AIEB)	Tr
Emotional and Psychological	Served as the primary source of emotional regulation, daily venting, and deep vulnerability.[27]	Th th so sy
Physical and Spatial	Shared living spaces, frequent physical affection, cuddling, and co-sleeping.[27]	ar dl ph
Temporal and Communication	Constant text messaging, expectation of immediate replies, shared daily routines.[29]	tic at lir pr

Medical and Caregiving	Attending psychiatric appointments together, actively monitoring medication compliance, managing side effects.[30]	wl ba or

Setting these boundaries often induces profound guilt in the healthy partner, who may feel they are selfishly abandoning a sick loved one during their darkest hour.[21] However, clinical recovery frameworks emphasize that establishing boundaries is a required form of self-care, not a punitive measure.[7] The healthy partner must recognize and internalize the "law of relationships": an individual is only responsible for their own thoughts, feelings, and actions, and cannot take responsibility for the neurochemistry or emotional states of another adult.[32]

Navigating the Guilt of De-escalation

The healthy partner must actively combat the societal and internal narrative that distancing themselves equates to abandonment. Threats of self-harm, deep despair, or rapid shifts between emotional extremes from the depressed partner must be handled by trained medical professionals, not the ex-partner.[21]

The transition requires a fundamental shift from active caregiving to passive support. Validating the depressed partner's experience without attempting to fix it is the cornerstone of this new dynamic. Utilizing phrases such as, "I can see how hard this medication adjustment is for you," replaces actionable problem-solving and protects the healthy partner from

over-investing.[31] The healthy partner must completely relinquish the fantasy that their enduring patience will miraculously cure the AIEB.

Medical Interventions for Antidepressant-Induced Emotional Blunting

While the non-depressed partner focuses entirely on boundary enforcement and self-preservation, the ultimate trajectory of the friendship is heavily dependent on whether the depressed partner takes independent, clinical action to address their AIEB. The emotional withdrawal is not a permanent, organic reality; it is a highly manageable pharmacological side effect that responds well to psychiatric intervention.[4]

When AIEB threatens to destroy significant interpersonal bonds, psychiatric guidelines recommend a proactive, systematic, stepwise approach to medication management. Routine screening utilizing tools like the Oxford Depression Questionnaire (ODQ) is recommended to quantify the severity of the blunting.[8]

1. Dose Reduction (First-Line Strategy)

Because AIEB is widely recognized as a dose-dependent side effect, the primary clinical recommendation is to reduce the current antidepressant dose by 25% to 50%, provided it is clinically feasible and does not risk a severe depressive relapse.[8] Dose changes require careful monitoring every 2 to 4 weeks. In ambiguous cases where it is unclear if the symptoms represent residual depression or a medication side effect,

a prescriber may utilize a brief 24-to-48-hour "drug holiday," or paradoxically, a brief dose increase. If the emotional blunting worsens during a dose increase, this definitively confirms a drug-induced etiology and supports subsequent dose reduction protocols.[8]

2. Switching to Lower-Risk Pharmacological Agents

If dose reduction results in the rapid return of severe depressive symptoms, the prescriber will typically switch the patient to a non-serotonergic or multimodal agent.[8]

Pharmacological Agent	Mechanism of Action	Bl
Bupropion	Norepinephrine-Dopamine Reuptake Inhibitor (NDRI). Avoids serotonin pathways entirely.	qu en m tic re
Vortioxetine	Multimodal serotonergic agent (modulates various 5-HT receptors directly).	st m a wi

Agomelatine / Mirtazapine	Melatonergic agonist / 5-HT2C receptor antagonist.	bl do pr in R
Alternative SSRIs	Selective Serotonin Reuptake Inhibitors (e.g., transitioning from Sertraline to Fluoxetine).	th de m to fo ca

3. Pharmacological Augmentation Strategies

For patients whose clinical depression is uniquely and successfully controlled by an SSRI, discontinuing the medication may present too high a risk of psychiatric relapse. In these instances, adding a secondary agent can act as a targeted "antidote" to the blunting.[1] Adding a low dose of bupropion (e.g., 150 mg/day SR/XL) can directly counteract the serotonin-induced dopamine suppression in the reward pathways, restoring emotional engagement and vitality without compromising the primary antidepressant effect.[8]

4. Alternative Neuromodulation Therapies

If oral pharmacological interventions fail to adequately balance mood stabilization with emotional preservation, patients may explore non-systemic, alternative treatments. Deep

Transcranial Magnetic Stimulation (TMS) is a non-invasive, FDA-approved procedure that uses magnetic pulses to regulate specific mood circuitry in the brain. Because it does not introduce systemic chemical suppression like oral antidepressants, it allows patients to achieve remission from depression while maintaining their full emotional range and capacity for connection.[2]

The non-depressed partner's role regarding these medical interventions must remain strictly observational. They can provide specific feedback on the behavioral changes they have observed and suggest the patient speak to a psychiatrist, but they must entirely recuse themselves from taking responsibility for the medication management or acting as the patient's medical advocate.[2]

Ecosystems of Care: Utilizing Localized Support Frameworks

Successfully navigating the psychological fallout of an SSRI-induced relationship transition requires accessing specialized professional support. Individuals cannot process this level of ambiguous loss and psychiatric complexity in a vacuum. For both the partner experiencing AIEB and the healthy partner seeking to establish boundaries, local ecosystems of care provide vital infrastructure. Analyzing a specific regional context—such as the Magic Valley area of Idaho, encompassing Twin Falls and Buhl—illustrates exactly how these clinical resources can be mobilized to facilitate healthy transitions.[35]

Specialized Therapeutic Interventions for Relationship Transitions

For the non-depressed partner processing the ambiguous loss and trauma of the transition, securing individual therapy is paramount to prevent long-term attachment issues.[1] In the Buhl and Twin Falls area, several specific practices offer highly targeted modalities for this exact demographic:

- **A Chance 4 Change (Twin Falls):** Located on Shoshone Street, this clinic specializes heavily in life transitions, relationship distress, and complex family dynamics.[35] Clinicians such as Maira Torres, LMSW, and Evan Maynard, LMSW, focus specifically on navigating life transitions, which is crucial for partners adjusting to the abrupt end of a romantic bond.[35] The clinic utilizes Solution Focused Brief Therapy (SFBT) and Cognitive Behavioral Therapy (CBT), methods empirically proven to be effective in establishing firm boundaries, processing grief, and rebuilding self-concept after caregiver burnout.[2]
- **Trena H. Peckham, LMFT (Buhl):** Specializing in Emotionally Focused Therapy (EFT) and Internal Family Systems (IFS), this practice addresses deep attachment wounds and severe relationship distress. IFS provides an excellent framework to help individuals untangle their core identity from the "rescuer" or caregiving role they assumed during their partner's depression.[38]
- **Paris Conwell, LPC (Buhl):** Focuses heavily on anxiety, life transitions, and codependency utilizing

CBT, experiential, and somatic approaches. This provides the necessary clinical interventions for healthy partners struggling with the pervasive anxiety of keeping a relationship in an undefined limbo.[40]

- **Turning Point Center for Change (Twin Falls):** Offers comprehensive behavioral health services, including evidence-based modalities like EMDR and ART, which are highly effective for processing the trauma associated with sudden relationship dissolution and emotional abandonment.[42]

Psychiatric and Medical Management Resources

For the partner actively experiencing AIEB, localized medical intervention is absolutely required to adjust SSRI protocols, pursue augmentation, or explore alternatives:

- **Family Health Services (Buhl & Twin Falls):** Offers highly integrated Primary Care Behavioral Health and specialized psychiatric medication management. This infrastructure allows the patient to address the pharmacological origins of their blunting alongside therapeutic support, utilizing sliding fee scales to remove financial barriers to care.[36]
- **Magic Valley Behavioral Health (Twin Falls):** Led by Dr. Kelsie Hendrickson, this clinic offers evidence-based treatments and consultations for complex psychiatric presentations, ensuring that delicate medication adjustments and switching strategies are han-

dled by specialized professionals rather than general practitioners.[42]

Community Support and Crisis Networks

The transition from romantic partner to platonic friend often leaves both individuals severely isolated, having lost their primary source of social interaction. Community support groups mitigate this isolation by connecting individuals with peers experiencing remarkably similar relational dynamics.[42]

- **NAMI (National Alliance on Mental Illness):** The NAMI Idaho chapters offer robust, peer-led support systems accessible to Magic Valley residents.[46] The **NAMI Family Support Group** is specifically designed for loved ones and caregivers of individuals with mental illness. This provides a critical, confidential space for the non-depressed partner to gain insight, process caregiver burnout, and learn boundary-setting techniques from others facing similar circumstances.[46] Concurrently, the **NAMI Connection Recovery Support Group** serves the individual actively navigating the psychiatric diagnosis.[42]
- **Crisis Center of South Central Idaho:** For acute mental health emergencies that may arise during the volatile de-escalation of the relationship, this Twin Falls facility provides emergency behavioral health services without charge for up to 23 hours. This completely removes the burden of crisis management from the ex-partner.[47]

Strategic Decision Framework: Sustaining Friendship versus Executing a Clean Break

For the non-depressed partner, navigating the enduring "limbo" of this transition requires a brutal, objective assessment of their own mental health and the depressed partner's willingness to engage in clinical recovery. The ultimate decision to sustain a platonic friendship or execute a clean break hinges on several critical, observable factors.

Clinical Indicators Supporting a Transition to Friendship

Remaining friends may be a viable, non-destructive path if, and only if, the following conditions are consistently met:

1. **Clinical Insight and Accountability:** The depressed partner possesses deep insight into their condition, openly acknowledging that their emotional withdrawal and apathy are linked directly to their SSRI treatment, rather than an inherent flaw in the healthy partner or the relationship.[2]
2. **Active Medical Management:** The depressed partner does not accept the blunting as a permanent state. They are actively collaborating with their prescriber to adjust doses, switch medications to lower-risk agents like bupropion, or pursue therapeutic augmentation strategies.[2]
3. **Strict Respect for Boundaries:** The depressed partner honors the mandated no-contact period and fully adheres to the new, platonic parameters of the relationship without resorting to emotional manipu-

lation, guilt-tripping, or demanding unauthorized caregiver labor.[48]

4. **Mutual Independence and Self-Expansion:** The healthy partner has successfully divested their emotional dependency on the depressed partner. They have actively re-established their own support networks, re-engaged in personal hobbies, and restored their self-esteem entirely independent of the ex-partner's validation.[26]

Clinical Indicators Necessitating a Clean Break

Conversely, attempting to maintain a friendship will invariably result in chronic psychological harm to the healthy partner if the following toxic dynamics are present:

1. **Persistent Denial and Blame:** The depressed partner denies the systemic impact of the medication, projecting the relationship's failure entirely onto the healthy partner or insisting that their chemically blunted state represents their "true," objective feelings.[28]
2. **Refusal to Adjust Psychiatric Treatment:** The depressed partner vehemently refuses to discuss side effects with a physician, preferring the comfortable safety of emotional numbness over the necessary vulnerability of emotional recovery and medication adjustment.[2]
3. **Egregious Boundary Violations:** The depressed partner continually expects the healthy partner to function as a therapist, crisis manager, or primary emotional regulator. They effectively demand all the supportive

benefits of a dedicated romantic relationship without offering any reciprocal affection, commitment, or intimacy.[28]

4. **Severe Caregiver Burnout:** The healthy partner experiences severe sleep disturbances, persistent anxiety, deep-seated resentment, or a profound loss of personal identity that is directly tied to monitoring, worrying about, or managing the depressed partner's life.[18]

If the relationship must end entirely, the healthy partner must prioritize radical self-care. They must acknowledge the clinical reality that one cannot cure another person's neurochemistry through sheer willpower, enduring presence, or self-sacrifice.[12]

The request to "stay friends" while one partner undergoes an emotionally blunting SSRI treatment creates a highly volatile, asymmetrical interpersonal dynamic. Antidepressant-Induced Emotional Blunting is a profound neurochemical alteration that shifts a patient from affective engagement to a state of complete cognitive detachment.[8] While the medication successfully dampens the acute agony of clinical depression, it inadvertently starves the dopamine-driven reward pathways entirely necessary for maintaining romantic passion, joy, and relational empathy.[1]

For the non-depressed partner, blindly accommodating this transition without strict boundaries is psychologically hazardous. It precipitates a state of ambiguous loss, wherein the healthy partner acts as an emotional regulator and unpaid caregiver while being denied the reciprocal intimacy of a relationship.[5] To successfully navigate this crisis, the healthy part-

ner must execute a strategic withdrawal, enforce rigid boundaries, and utilize specialized therapeutic resources to reclaim their emotional autonomy. The preservation of one's mental health must always supersede the futile desire to rescue a partner from the side effects of their own psychopharmacology.

Works cited

1. SSRI Emotional Blunting: Understanding, Recognizing, and Restoring Your Emotional Life on Antidepressants - Dr. Beata Lewis, MD, accessed April 20, 2026, https://drlewis.com/ssri-emotional-blunting-complete-guide/
2. Emotional Blunting from Antidepressants: What to Do, accessed April 20, 2026, https://axismh.com/emotional-blunting-from-antidepressants/
3. Scientists explain emotional 'blunting' caused by common antidepressants, accessed April 20, 2026, https://www.cam.ac.uk/research/news/scientists-explain-emotional-blunting-caused-by-common-antidepressants
4. Can SSRIs Make Me Fall Out Of Love? - Neuro Wellness Spa, accessed April 20, 2026, https://neurowellnessspa.com/can-antidepressants-make-me-fall-out-of-love/
5. How Emotional Numbness Steals Your Relationship | Psychology ..., accessed April 20, 2026, https://www.psychologytoday.com/us/blog/the-an-

gry-therapist/202503/how-emotional-numbness-steals-your-relationship

6. Supporting partners through mental health recovery: tackling caregiver burnout - Orchestrate Healthcare, accessed April 20, 2026, https://www.orchestratehealth.com/supporting-partners-through-mental-health-recovery-tackling-caregiver-burnout/
7. The Critical Role of Boundaries for Family Caregivers - Psychology Today, accessed April 20, 2026, https://www.psychologytoday.com/us/blog/gaining-and-sustaining/202508/the-critical-role-of-boundaries-for-family-caregivers
8. Antidepressant-Induced Emotional Blunting: Diagnosis ..., accessed April 20, 2026, https://psychopharmacologyinstitute.com/publication/antidepressant-induced-emotional-blunting-diagnosis-mechanisms-and-management-2/
9. SSRI-Induced Indifference - PMC, accessed April 20, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC2989833/
10. Depression: Supporting a family member or friend - Mayo Clinic, accessed April 20, 2026, https://www.mayoclinic.org/diseases-conditions/depression/in-depth/depression/art-20045943
11. Emotional Blunting, Cognitive Impairment, Bone Fractures, and Bleeding as Possible Side Effects of Long-Term Use of SSRIs - PMC, accessed April 20, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC8650205/

12. Protecting Relationships from Depression - Promises Behavioral Health, accessed April 20, 2026, https://www.promises.com/addiction-blog/relationships-depression/
13. Effect of Spousal Loss on Depression in Older Adults: Impacts of Time Passing, Living Arrangement, and Spouse's Health Status before Death - PMC, accessed April 20, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC8700949/
14. accessed April 20, 2026, https://www.betterhelp.com/advice/relations/feeling-numb-in-a-relationship-a-comprehensive-guide-to-healing-and-reconnection/#:~:text=Even%20healthy%20relationships%20can%20experience,isolation%2C%20loneliness%2C%20and%20resentment.
15. Supporting a partner with depression | Mental Health Foundation, accessed April 20, 2026, https://www.mentalhealth.org.uk/explore-mental-health/articles/supporting-partner-depression
16. Romantic Relationships and Mental Health: Investigating the Role of Self-Expansion on Depression Symptoms - PMC, accessed April 20, 2026, https://pmc.ncbi.nlm.nih.gov/articles/PMC12668558/
17. THE PSYCHOLOGICAL IMPACT ON THE RELATIONAL PARTNER OF BPD- DIAGNOSED PERSONS by Carl Pendleton Liberty University A Dissertatio, accessed April 20, 2026, https://digitalcommons.liberty.edu/cgi/viewcontent.cgi?article=6187&context=doctoral

18. Maintaining Boundaries As A Caregiver: Go From Guilt To Glow | Mental Health America, accessed April 20, 2026, https://mhanational.org/resources/maintaining-boundaries-as-a-caregiver-go-from-guilt-to-glow/
19. When Caregiving Affects Your Relationships - Benjamin Rose, accessed April 20, 2026, https://www.benrose.org/resource/when-caregiving-affects-your-relationships/
20. Intimacy when caring for your spouse - Caregiver Action Network, accessed April 20, 2026, https://www.caregiveraction.org/intimacy-when-caregiving-for-your-spouse/
21. Depression in Relationships: When to Say Goodbye - Healthline, accessed April 20, 2026, https://www.healthline.com/health/depression/relationships
22. How to take care of yourself after a breakup | Division of Student Life, accessed April 20, 2026, https://www.colorado.edu/studentlife/breakup-tips
23. Setting Boundaries in Recovery - Hazelden Betty Ford Foundation, accessed April 20, 2026, https://www.hazeldenbettyford.org/articles/boundaries-in-addiction-recovery
24. What do healthy boundaries look like in recovery? - MHA Screening, accessed April 20, 2026, https://screening.mhanational.org/content/what-do-healthy-boundaries-look-like-in-recovery/
25. How to Manage Anxiety After a Breakup | Coping Strategies - Baltimore Therapy Group, accessed April

20, 2026, https://www.therapistsinbaltimore.com/therapy-insights-blog/2025/12/6/how-to-manage-anxiety-after-a-breakup-coping-strategies
26. 6 Ways to Maintain Your Mental Health During a Break-Up - One Love Foundation, accessed April 20, 2026, https://www.joinonelove.org/learn/6-ways-to-maintain-your-mental-health-during-a-break-up/
27. Setting Healthy Boundaries in Relationships - HelpGuide.org, accessed April 20, 2026, https://www.helpguide.org/relationships/social-connection/setting-healthy-boundaries-in-relationships
28. relationship, splitting, emotional regulation : r/BPD - Reddit, accessed April 20, 2026, https://www.reddit.com/r/BPD/comments/1spjt21/relationship_splitting_emotional_regulation/
29. Setting Boundaries in Healthy Relationships - BHSI, accessed April 20, 2026, https://bhsiclinics.com/setting-boundaries-in-healthy-relationships/
30. 7 Ways to Help your Partner Deal with Depression - HealthCentral, accessed April 20, 2026, https://www.healthcentral.com/condition/depression/how-to-help-your-partner-with-depression
31. 7 Ways To Support Your Partner's Mental Health (When You're Not a Therapist), accessed April 20, 2026, https://mentalhealthfirstaid.org/news/7-ways-to-support-your-partners-mental-health/
32. Map it out: Setting boundaries for your well-being - Mayo Clinic Health System, accessed April 20, 2026, https://www.mayoclinichealthsystem.org/hometown-

health/speaking-of-health/setting-boundaries-for-well-being
33. Dealing with BPD in Relationships Tips - HelpGuide.org, accessed April 20, 2026, https://www.helpguide.org/mental-health/personality-disorders/bpd-in-relationships
34. Dos and Don'ts of Supporting a Partner With Depression - Everyday Health, accessed April 20, 2026, https://www.everydayhealth.com/mental-health-conditions/dos-and-donts-of-supporting-partner-with-depression/
35. Melissa Osen - Behavioral health therapy, accessed April 20, 2026, https://achance4change.clientsecure.me/
36. Behavioral Health Services - Therapists | Family Health Services, accessed April 20, 2026, https://fhsid.org/services/behavioral-health/
37. Mental Health Treatment in Twin Falls, Idaho - The Walker Center, accessed April 20, 2026, https://www.thewalkercenter.org/mental-health-services
38. About Trena Peckman | Trauma-Informed Therapy | Idaho & Alabama, USA, accessed April 20, 2026, https://trenapeckham.com/trena-peckham
39. Trena H Peckham, Marriage & Family Therapist, Buhl, ID, 83316 | Psychology Today, accessed April 20, 2026, https://www.psychologytoday.com/us/therapists/trena-h-peckham-buhl-id/349188
40. Therapist Paris - Counseling, accessed April 20, 2026, https://therapist-paris.clientsecure.me/

41. Paris Conwell, Licensed Professional Counselor, Buhl, ID, 83316 | Psychology Today, accessed April 20, 2026, https://www.psychologytoday.com/us/therapists/paris-conwell-buhl-id/1450281
42. Mental Health Resources in Twin Falls: A Complete Guide to Local Support Services, accessed April 20, 2026, https://tpcfc.org/mental-health-resources-in-twin-falls-a-complete-guide-to-local-support-services/
43. Turning Point Center for Change: Counselors in Twin Falls, ID, accessed April 20, 2026, https://tpcfc.org/
44. Magic Valley Behavioral Health: Psychologist in Twin Falls, Idaho | Kelsie Hendrickson, PhD, ABPP, accessed April 20, 2026, https://www.mvbh.org/
45. Navigating Relationships in Recovery: Love, Friendship & Boundaries, accessed April 20, 2026, https://guidingroad.com/recovery-journal/navigating-relationships-in-recovery-love-friendship-boundaries/
46. Support Groups - NAMI Idaho, accessed April 20, 2026, https://namiidaho.org/support-and-education/support-groups/
47. Mental Health & Wellness Resources - LINC Idaho, accessed April 20, 2026, https://www.lincidaho.org/health-wellness
48. The Scope Blog - Navigating Relationships After Mental Health Recovery, accessed April 20, 2026, https://www.tanner.org/the-scope/navigating-relationships-after-mental-health-recovery

49. Ways to Build Healthy Relationships in Recovery | The Pavilion at Williamsburg Place, accessed April 20, 2026, https://pavilionwp.com/ways-to-build-healthy-relationships-in-recovery/
50. Depressed Spouse: How to Help Your Partner - WebMD, accessed April 20, 2026, https://www.webmd.com/depression/features/help-depressed-spouse